

Energy of commuting graph of finite groups whose centralizers are Abelian

Reza Sharafadini ^{*}, Rezvan Darbandi

Abstract. Let G be a finite group with centre $Z(G)$. The commuting graph of a non-Abelian group G , denoted by Γ_G , is a simple undirected graph whose vertex set is $G \setminus Z(G)$, and two vertices x and y are adjacent if and only if $xy = yx$. In this article we aim to compute the ordinary energy of Γ_G for groups G whose centralizers are Abelian.

1. Introduction

Let G be a finite group with centre $Z(G)$. The commuting graph of a non-Abelian group G , denoted by Γ_G , is a simple undirected graph whose vertex set is $G \setminus Z(G)$, and two vertices x and y are adjacent if and only if $xy = yx$. Various properties of commuting graphs of finite groups have been studied (see [4, 6, 16, 17, 19, 21, 23]). Note that the complement of Γ_G is called non-commuting graph of G (see [1, 18]). Indeed, for a non-Abelian group G , the *non-commuting graph* $\overline{\Gamma}_G$ of G has the vertex set $G \setminus Z(G)$ and two distinct vertices x and y are adjacent if $xy \neq yx$. According to [20] non-commuting graphs were first considered by Erdős in 1975. Over the past decade, non-commuting graphs have received considerable attention. For example, Abdollahi et al. [1] proved that the diameter of any non-commuting graph is 2. For two non-Abelian groups with isomorphic non-commuting graphs, the sufficient conditions that guarantee their orders are equal were provided by Abdollahi and Shahverdi [2] and Darafsheh [9]. Akbari and Moghaddamfar [5] studied strongly regular non-commuting

2000 Mathematics Subject Classification: 05C12, 05C25, 20D60, 05A15.

Keywords: Commuting graph, non-commuting graph, AC-group, Eigenvalue, Energy.

^{*} Corresponding author

graphs. Solomon and Woldar [23] characterized some simple groups by their non-commuting graphs.

For any element x of a group G , the set $C_G(x) = \{y \in G \mid xy = yx\}$ is called the centralizer of x in G . Let $|\text{Cent}(G)| = |\{C_G(x) \mid x \in G\}|$, that there is the number of distinct centralizers in G . A group G is called an n -centralizer group if $|\text{Cent}(G)| = n$. In [8], Belcastro and Sherman characterized finite n -centralizer groups for $n = 4, 5$ and they showed that there is no finite n -centralizer group for $n \in \{2, 3\}$ (while Ashrafi in [3], showed that, for any positive integer $n \neq 2, 3$, there exists a finite group G such that $|\text{Cent}(G)| = n$). A group G is called an AC-group if $C_G(x)$ is Abelian for every $x \in G \setminus Z(G)$. Various aspects of AC-groups can be found in [1, 10, 22].

Let Γ be a graph with the adjacency matrix A . Let $\lambda_1, \lambda_2, \dots, \lambda_t$ be the distinct eigenvalues of A with corresponding mutiplicites $m_1, m_2, \dots, m_t$. The spectrum of Γ is defined as

$$\text{Spec}(\Gamma) = \{\lambda_1^{(m_1)}, \lambda_2^{(m_2)}, \dots, \lambda_t^{(m_t)}\}.$$

The energy of graph Γ is defined as the sum of absolute values of its eigenvalues [13, 15],

$$\mathcal{E}(\Gamma) = \sum_{i=1}^t m_i |\lambda_i|.$$

The energy is a graph parameter stemming from the Hückel molecular orbital approximation for the total π -electron energy (for recent survey on molecular graph energy see [14] and [7]).

Let Γ_1 and Γ_2 be two graphs with disjoint vertex sets V_1 and V_2 . The union of Γ_1 and Γ_2 , denoted by $\Gamma_1 \sqcup \Gamma_2$ is the graph whose vertex set and edge set are $V_1 \cup V_2$ and $E(\Gamma) = E_1 \cup E_2$, respectively.

The complete graph on n vertices is denoted by K_n . It is well known that $\text{Spec}(K_n) = \{(-1)^{(n-1)}, (n-1)^1\}$. Further, if $\Gamma = K_{n_1} \sqcup K_{n_2} \sqcup \dots \sqcup K_{n_l}$, then

$$\text{Spec}(\Gamma) = \left\{ (-1)^{\sum_{i=1}^l n_i - l}, (n_1 - 1)^1, (n_2 - 1)^1, \dots, (n_l - 1)^1 \right\}. \quad (1)$$

2. Main Results

In this section, we recall the spectrum of the commuting graphs of some particular families of AC-groups (see [11, 12]). Then we compute their

energies.

Theorem 1 ([18, Proposition 3.1]). *Let G be a finite non-Abelian AC-group. Then the commuting graph $\Gamma(G)$ is a union of complete graphs if and only if G is an AC-group.*

Let $C_1, \dots, C_n$ be the distinct centralizers of non-central elements of G . Then it follows from the proof of Theorem 1 that $C_i \cap C_j = Z(G)$ for all $1 \leq i \neq j \leq n$. It follows that $\Gamma(G) = \bigsqcup_{i=1}^n K_{|C_i| - |Z(G)|}$. It follows that the following theorem is a consequence of Theorem 1.

Theorem 2 ([12]). *Let G be a finite non-Abelian AC-group. Let $C_1, \dots, C_n$ be the distinct centralizers of non-central elements of G . Then*

$$\text{Spec}(\Gamma_G) = \left\{ (-1)^{\sum_{i=1}^n |C_i| - n(|Z(G)|+1)}, (|C_1| - |Z(G)| - 1)^1, \dots, (|C_n| - |Z(G)| - 1)^1 \right\}$$

Lemma 1. *Let G be a finite non-Abelian group. Let $C_1, \dots, C_n$ be the centralizers of non-central elements of G . Then*

$$\sum_{i=1}^n |C_i| = |G|(k(G) - |Z(G)|),$$

where $k(G)$ denotes the number of conjugacy classes of G .

Proof. Let $g_1, \dots, g_{k(G)}$ be the representative of conjugacy classes of G and without loss of generality assume that $g_1, \dots, g_t$ are non-central. In this case, $k(G) = |Z(G)| + t$. Let us denote the conjugacy class of G corresponding to g_i , by $\overline{g_i}$. It follows from the fact $|\overline{g_i}| = \frac{|G|}{|C_G(g_i)|}$ that $|C_G(g)| = |C_G(g_i)|$ for all $g \in \overline{g_i}$. If $g \notin Z(G)$, then $g \in \overline{g_i}$ for some $1 \leq i \leq t$. It follows that

$$\sum_{g \notin Z(G)} |C_G(g)| = \sum_{i=1}^t \sum_{g \in \overline{g_i}} |C_G(g_i)| = \sum_{i=1}^t |\overline{g_i}| |C_G(g_i)| = t|G| = |G|(k(G) - |Z(G)|).$$

□

The following theorem is a direct consequence of Theorem 2 and Lemma 1

Theorem 3. *Let G be a finite non-Abelian AC-group with n distinct centralizers of non-central elements. Then*

$$\mathcal{E}(\Gamma_G) = 2|G|(k(G) - |Z(G)|) - 2n(|Z(G)| + 1),$$

where $k(G)$ denotes the number of conjugacy classes of G .

Corollary 1 ([12]). *Let G be a finite non-Abelian AC-group and A be any finite Abelian group. Then the spectrum of the commuting graph of $G \times A$ is given by*

$$\left\{ (-1)^{\sum_{i=1}^n |A|(|C_i| - n|Z(G)|) - n}, (|A|(|C_1| - |Z(G)|) - 1)^1, \dots, (|A|(|C_n| - |Z(G)|) - 1)^1 \right\}$$

where $C_1, \dots, C_n$ are the distinct centralizers of non-central elements of G .

Lemma 2 ([12]). (i) *Let $M_{2mn} = \langle a, b : a^m = b^{2n} = 1, bab^{-1} = a^{-1} \rangle$ be a metacyclic group, where $m > 2$. Then*

$$\text{Spec}(\Gamma_{M_{2mn}}) = \begin{cases} \left\{ (-1)^{2mn-m-n-1}, (n-1)^m, (mn-n-1)^1 \right\} & \text{if } m \text{ is odd} \\ \left\{ (-1)^{2mn-2n-\frac{m}{2}-1}, (2n-1)^{\frac{m}{2}}, (mn-2n-1)^1 \right\} & \text{if } m \text{ is even.} \end{cases}$$

(ii)

$$\text{Spec}(\Gamma_{D_{2m}}) = \begin{cases} \{ (-1)^{m-2}, 0^m, (m-2)^1 \} & \text{if } m \text{ is odd} \\ \{ (-1)^{\frac{3m}{2}-3}, 1^{\frac{m}{2}}, (m-3)^1 \} & \text{if } m \text{ is even.} \end{cases}$$

(iii) *The spectrum of the commuting graph of the dicyclic group or the generalized quaternion group*

$$Q_{4m} = \langle a, b : a^{2m} = 1, b^2 = a^m, bab^{-1} = a^{-1} \rangle,$$

where $m \geq 2$, is given by

$$\text{Spec}(\Gamma_{Q_{4m}}) = \{ (-1)^{3m-3}, 1^m, (2m-3)^1 \}.$$

(iv) *Consider the group $U_{6n} = \{a, b : a^{2n} = b^3 = 1, a^{-1}ba = b^{-1}\}$. Then*

$$\text{Spec}(\Gamma_{U_{6n}}) = \{ (-1)^{5n-4}, (n-1)^3, (2n-1)^1 \}.$$

Corollary 2.

$$\mathcal{E}(\Gamma_{M_{2mn}}) = \begin{cases} 4mn - 2m - 2n - 2 & \text{if } m \text{ is odd} \\ 4mn - 4n - m - 2 & \text{if } m \text{ is even.} \end{cases}$$

$$\mathcal{E}(\Gamma_{D_{2m}}) = \begin{cases} 2m - 4 & \text{if } m \text{ is odd} \\ 3m - 6 & \text{if } m \text{ is even.} \end{cases}$$

$$\mathcal{E}(\Gamma_{Q_{4m}}) = 6m - 3.$$

$$\mathcal{E}(\Gamma_{U_{6n}}) = 10n - 8.$$

Lemma 3 ([11]). *The spectrum of the commuting graph of the quasidihedral group*

$$QD_{2^n} = \langle a, b \mid a^{2^{n-1}} = b^2 = 1, bab^{-1} = a^{2^{n-2}-1} \rangle,$$

where $n \geq 4$, is given by

$$\text{Spec}(\Gamma_{QD_{2^n}}) = \left\{ (-1)^{2^n-2^{n-2}-3}, 1^{2^{n-2}}, (2^{n-1}-3)^1 \right\}.$$

Lemma 4 ([11]). *The spectrum of the commuting graph of the projective special linear group $PSL(2, 2^k)$, where $k \geq 2$, is given by*

$$\text{Spec}(\Gamma_{PSL(2, 2^k)}) = \left\{ (-1)^{2^{3k}-2^{2k}-2^{k+1}-2}, (2^k-1)^{2^{k-1}(2^k-1)}, (2^k-2)^{2^k+1}, (2^k-3)^{2^{k-1}(2^k+1)} \right\}.$$

Lemma 5 ([11]). *The spectrum of the commuting graph of the general linear group $GL(2, q)$, where $q = p^n > 2$ and p is a prime integer, is given by*

$$\text{Spec}(\Gamma_{GL(2, q)}) = \left\{ (-1)^{q^4-q^3-2q^2-q}, (q^2-3q+1)^{\frac{q(q+1)}{2}}, (q^2-q-1)^{\frac{q(q-1)}{2}}, (q^2-2q)^{q+1} \right\}.$$

Theorem 4 ([11]). *Let G be a finite group and $\frac{G}{Z(G)} \cong Sz(2)$, where*

$$Sz(2) = \langle a, b \mid a^5 = b^4 = 1, b^{-1}ab = a^2 \rangle.$$

Then

$$\text{Spec}(\Gamma_G) = \left\{ (-1)^{(19|Z(G)|-6)}, (4|Z(G)|-1)^1, (3|Z(G)|-1)^5 \right\}.$$

Lemma 6 ([11]). *Let $F = GF(2^n)$, $n \geq 2$ and ϑ be the Frobenius automorphism of F , i. e., $\vartheta(x) = x^2$ for all $x \in F$. Then the spectrum of the commuting graph of the group*

$$A(n, \vartheta) = \left\{ U(a, b) = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & \vartheta(a) & 1 \end{bmatrix} \mid a, b \in F \right\}.$$

under matrix multiplication given by $U(a, b)U(a', b') = U(a + a', b + b' + a'\vartheta(a))$ is

$$\text{Spec}(\Gamma_{A(n, \vartheta)}) = \left\{ (-1)^{(2^n-1)^2}, (2^n - 1)^{2^n-1} \right\}.$$

Lemma 7 ([11]). *Let $F = GF(p^n)$, p be a prime. Then the spectrum of the commuting graph of the group*

$$A(n, p) = \left\{ V(a, b, c) = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & c & 1 \end{bmatrix} \mid a, b, c \in F \right\}.$$

under matrix multiplication $V(a, b, c)V(a', b', c') = V(a+a', b+b'+ca', c+c')$ is

$$\text{Spec}(\Gamma_{A(n, p)}) = \left\{ (-1)^{p^{3n}-2p^n-1}, (p^{2n} - p^n - 1)^{p^n+1} \right\}.$$

Lemma 8 ([11]). *Let G be a non-Abelian group of order pq , where p and q are primes with $p \mid (q-1)$. Then*

$$\text{Spec}(\Gamma_G) = \{(-1)^{pq-q-1}, (p-2)^q, (q-2)^1\}.$$

Corollary 3. (i)

$$\mathcal{E}(\Gamma_{QD_{2^n}}) = 2^n + 2^{n-1} - 6.$$

(ii)

$$\begin{aligned} \mathcal{E}(\Gamma_{PSL(2, 2^k)}) &= (2^{3k} - 2^{2k} - 2^{k+1} - 2) + (2^k - 1)(2^{k-1}(2^k - 1)) \\ &\quad + (2^k - 2)(2^k + 1) + (2^k - 3)(2^{k-1}(2^k + 1)). \end{aligned}$$

(iii)

$$\begin{aligned} \mathcal{E}(\Gamma_{GL(2, q)}) &= (q^4 - q^3 - 2q^2 - q) + (q^2 - 3q + 1)\left(\frac{q(q+1)}{2}\right) \\ &\quad + (q^2 - q - 1)\left(\frac{q(q-1)}{2}\right) + (q^2 - 2q)(q+1). \end{aligned}$$

(iv) Let G be a finite group and $\frac{G}{Z(G)} \cong Sz(2)$. Then

$$\mathcal{E}(\Gamma_G) = 28|Z(G)| - 12.$$

(v)

$$\mathcal{E}(\Gamma_{A(n,\vartheta)}) = (2^n - 1)^2 + (2^n - 1)(2^n - 1) = 2(2^n - 1)^2.$$

(vi)

$$\mathcal{E}(\Gamma_{A(n,p)}) = (p^{3n} - 2p^n - 1) + (p^n + 1)(p^{2n} - p^n - 1) = 2p^{3n} - 4p^n - 2.$$

(vii) Let G be a non-Abelian group of order pq , where p and q are primes with $p \mid (q - 1)$. Then

$$\mathcal{E}(\Gamma_G) = 2q(p - 1) - 3.$$

Note that $A_5 \cong PSL(2, 4)$. Then by Lemma 4, we have

$$\text{Spec}(\Gamma_{A_5}) = \{(-1)^{38}, 1^{10}, 2^5, 3^6\}, \quad \mathcal{E}(\Gamma_{A_5}) = 76.$$

Theorem 5 ([12]). Let G be a finite group, p be a prime and $m \geq 2$ a natural number. Then the following holds.

(i) If $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, then G is AC-group and

$$\text{Spec}(\Gamma_G) = \left\{ (-1)^{(p^2-1)|Z(G)|-p-1}, ((p-1)|Z(G)|-1)^{p+1} \right\}.$$

(ii) If $\frac{G}{Z(G)} \cong D_{2m}$, then G is AC-group and

$$\text{Spec}(\Gamma_G) = \left\{ (-1)^{(2m-1)|Z(G)|-m-1}, (|Z(G)|-1)^m, ((m-1)|Z(G)|-1)^1 \right\}.$$

Corollary 4. Let G be a finite group, p be a prime and $m \geq 2$ a natural number. Then the following hold.

(i) If $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, then $\mathcal{E}(\Gamma_G) = 2((p^2 - 1)|Z(G)| - p - 1)$.

(ii) If $\frac{G}{Z(G)} \cong D_{2m}$, then $\mathcal{E}(\Gamma_G) = 2((2m - 1)|Z(G)| - m - 1)$.

Corollary 5. Let G be a finite group and p be a prime. Then

- (i) Let G be a non-Abelian group of order p^3 , for any prime p , then $\mathcal{E}(\Gamma_G) = 2(p^3 - 2p - 1)$.
- (ii) If G is a finite 4-centralizer group, then $\mathcal{E}(\Gamma_G) = 6(|Z(G)| - 1)$.
- (iii) If G is a finite $(p+2)$ -centralizer p -group, for any prime p , then

$$\mathcal{E}(\Gamma_G) = 2((p^2 - 1)|Z(G)| - p - 1).$$

- (iv) If G is a finite 5-centralizer group, then

$$\mathcal{E}(\Gamma_G) \in \{8(2|Z(G)| - 1), 10|Z(G)| - 8\}.$$

Proof. (i) In this case $|Z(G)| = p$ and $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$. Hence the result follows from Corollary 4(i).

(ii) The result follows from Corollary 4(i), since if G is a finite 4-centralizer group, then it follows from [8, Theorem 2] that $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

(iii) Let G be a finite $(p+2)$ -centralizer p -group. Then $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, by [3, Lemma 2.7]. Hence the result follows from Corollary 4(i).

(iv) If G is a finite 5-centralizer group, then by [8, Theorem 4] we have $\frac{G}{Z(G)} \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ or D_6 . If $\frac{G}{Z(G)} \cong \mathbb{Z}_3 \times \mathbb{Z}_3$, then by Corollary 4(i) we have $\mathcal{E}(\Gamma_G) = 8(2|Z(G)| - 1)$. In the case that $\frac{G}{Z(G)} \cong D_6$, by Theorem 5 we have $\mathcal{E}(\Gamma_G) = 10|Z(G)| - 8$. $\square$

As an application of Theorem 5 we have the following.

Corollary 6 ([11]). *Let G be a group isomorphic to any of the following groups*

- (i) $\mathbb{Z}_2 \times D_8$
- (ii) $\mathbb{Z}_2 \times Q_8$
- (iii) $M_{16} = \langle a, b \mid a^8 = b^2 = 1, bab = a^5 \rangle$
- (iv) $\mathbb{Z}_4 \rtimes \mathbb{Z}_4 = \langle a, b \mid a^4 = b^4 = 1, bab^{-1} = a^{-1} \rangle$
- (v) $D_8 * \mathbb{Z}_4 = \langle a, b, c \mid a^4 = b^2 = c^2 = 1, ab = ba, ac = ca, bc = a^2cb \rangle$
- (vi) $SG(16, 3) = \langle a, b \mid a^4 = b^4 = 1, ab = b^{-1}a^{-1}, ab^{-1} = ba^{-1} \rangle$.

Then $\text{Spec}(\Gamma_G) = \{(-1)^9, 3^3\}$, $\mathcal{E}(\Gamma_G) = 18$.

Proof. If G is isomorphic to any of the above listed groups, then $|G| = 16$ and $|Z(G)| = 4$. Therefore, $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. Thus the result follows from Theorem 5. $\square$

References

- [1] **A. Abdollahi, S. Akbari and H. R. Maimani**, Non-commuting graph of a group, *J. Algebra*, **298**, (2006), 468–492 .
- [2] **A. Abdollahi and H. Shahverdi**, Non-commuting graphs of nilpotent groups, *Comm. Algebra*, **42**, (2014), 3944–3949.
- [3] **A. R. Ashrafi**, On finite groups with a given number of centralizers, *Algebra Colloq.* **7**, (2000), 139–146.
- [4] **S. Akbari, A. Mohammadian, H. Radjavi and P. Raja**, On the diameters of commuting graphs, *Linear Algebra Appl.*, **418**, (2006), 161–176.
- [5] **M. Akbari and A.R. Moghaddamfar**, The existence or nonexistence of non-commuting graphs with particular properties, *J. Algebra Appl.*, **13** (2014), 1350064.
- [6] **C. Bates, D. Bundy, S. Hart and P. Rowley**, A Note on Commuting Graphs for Symmetric Groups, *Electron. J. Combin.* **16**, (2009), 1–13.
- [7] **R. A. Brualdi**, *Energy of a Graph*, available from <http://www.public.iastate.edu/~lhogben/energyB.pdf>
- [8] **S. M. Belcastro and G. J. Sherman**, Counting centralizers in finite groups, *Math. Magazine*, **67**, (1994), no. 5, 366–374.
- [9] **M. R. Darafsheh**, Groups with the same non-commuting graph, *Discrete Appl. Math.* **157**, (2009), 833–837.
- [10] **A. K. Das and D. Nongsang**, On the genus of the commuting graphs of finite non-Abelian groups, preprint, arXiv:1311.6324v1.
- [11] **J. Dutta and R. Nath**, Spectrum of commuting graphs of some classes of finite groups, <https://arxiv.org/abs/1604.07133>.
- [12] **J. Dutta and R. Nath**, Finite groups whose commuting graphs are integral, <https://arxiv.org/abs/1604.05902v1>.
- [13] **I. Gutman**, *The energy of a graph*, Ber. Math. Stat. Sekt. Forschungszent. Graz. **103**, (1978), 1–22.
- [14] **I. Gutman**, *The energy of a graph: old and new results*, Algebraic Combinatorics and Applications, Springer Berlin, 2001, 196–211.
- [15] **X. Li, Y. Shi, I. Gutman**, Graph Energy, Springer, New York, 2012.

- [16] **A. Iranmanesh and A. Jafarzadeh**, Characterization of finite groups by their commuting graph, *Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis* **23**(1), (2007), 7–13.
- [17] **A. Iranmanesh and A. Jafarzadeh**, On the commuting graph associated with the symmetric and alternating groups, *J. Algebra Appl.* **7** (1),(2008), 129–146.
- [18] **M. Mirzargar and A. R. Ashrafi**, Some Distance-Based Topological Indices Of A Non-Commuting Graph , *Hacettepe Journal of Mathematics and Statistics* **41**(4), (2012), 515–526.
- [19] **G. L. Morgan and C. W. Parker**, The diameter of the commuting graph of a finite group with trivial center, *J. Algebra* **393**(1), (2013), 41–59.
- [20] **B. H. Neumann**, A problem of Paul Erdős on groups, *J. Aust. Math. Soc. Ser. A* **21**, (1976), 467–472.
- [21] **C. Parker**, The commuting graph of a soluble group, *Bull. London Math. Soc.* **45**(4), (2013), 839–848.
- [22] **D. M. Rocke**, p -groups with abelian centralizers, *Proc. London Math. Soc.* **30**(3), (1975), 55–75.
- [23] **R. Solomon and A. Woldar**, Simple groups are characterized by their non-commuting graph, *J. Group Theory*, **16**, (2013), 793–824.

Reza Sharafadini
 Department of Mathematics
 Faculty of Science, Persian Gulf University,
 Bushehr 75169-13817
 IRAN
 e-mail: sharafadini@pgu.ac.ir

Rezvan Darbandi
 Department of Mathematics and Computer Science
 Amirkabir University of Technology
 Tehran
 IRAN e-mail: rez1.darnadi@aut.ac.ir